

MORRISON & FOERSTER LLP

Attorneys at Law
755 Page Mill Road
Palo Alto, California 94304-1018
Telephone: (650) 813-5600
Facsimile: (650) 494-0792

To: Examiner Scott Houtteman
Group Art Unit 1634
U. S. Patent & Trademark Office

Facsimile: (703) 305-7939
Phone: (703) 308-3885

From: Sean M. Brennan

Date: November 12, 1999

We are transmitting a total of 7 pages (including this page).
Original or hard copy to follow if this box is checked ☐.

If you do not receive all pages, please call (650) 813-5741 as soon as possible.

Preparer of this slip has confirmed that facsimile number given is correct: 3248/39.917

This facsimile contains confidential information which may also be privileged. Unless you are the addressee (or authorized to receive for the addressee), you may not copy, use, or distribute it. If you have received it in error, please advise Morrison & Foerster LLP immediately by telephone or facsimile and return it promptly by mail.

Comments: INFORMAL COMMUNICATION

Please see attached in connection with our phone interview this afternoon regarding 08/484,340.

Methods in Enzymology

Volume 152

*Guide to Molecular Cloning
Techniques*

EDITED BY

Shelby L. Berger

NATIONAL CANCER INSTITUTE
NATIONAL INSTITUTES OF HEALTH
BETHESDA, MARYLAND

Alan R. Kimmel

NATIONAL INSTITUTE OF DIABETES AND DIGESTIVE AND KIDNEY DISEASES
NATIONAL INSTITUTES OF HEALTH
BETHESDA, MARYLAND

1987



ACADEMIC PRESS, INC.

Harcourt Brace Jovanovich, Publishers

Orlando San Diego New York Austin
Boston London Sydney Tokyo Toronto

[9] Nick Translation

By JUDY MEINKOTH and GEOFFREY M. WAHL

Nick translation, or more precisely nick translocation, is a specific procedure for incorporating radioactive nucleotides into double-stranded DNA. The method takes advantage of the ability of *Escherichia coli* DNA polymerase I to combine the sequential addition of nucleotide residues to the 3'-hydroxyl terminus of a nick [generated by pancreatic deoxyribonuclease (DNase) I] with the elimination of nucleotides from the adjacent 5'-phosphoryl terminus. Linear, supercoiled, nicked, or gapped circular double-stranded molecules can be labeled to specific activities $>10^8$ cpm/ μ g with deoxynucleotide 5'-[α - 32 P]triphosphates by this technique.^{1,2} Since the nicks are introduced at random sites in the duplex, the method generates a population of radioactive fragments which partially overlap each other. At saturating levels of nucleotide triphosphates the size of the fragments is determined by the DNase concentration. Fragments approximately 500–1500 nucleotides long produce optimal signal-to-noise ratios when hybridized to immobilized DNA or RNA (see this volume [45, 61]), presumably due to their ability to hybridize with each other in overlapping complementary regions to form "networks" or "hyperpolymers."² While experiments consistent with hyperpolymer formation of nick-translated probes have been reported,^{3,4} the reproducibility and extent of hyperpolymer formation seem to be difficult to obtain, probably because of the critical dependence on probe size.^{2,4} Longer probes have been correlated with higher backgrounds (G. Wahl, unpublished observations).

Nick Translation Reaction

1. Dilute stock DNase in 50 mM Tris-HCl at pH 7.4, 10 mM MgCl₂, 1 mg/ml DNase-free bovine serum albumin just before use. A DNase concentration of about 40–80 pg/ μ l is usually adequate.
2. Mix 2 μ l 10 \times NT (nick translation) buffer (10 \times = 50 mM MgCl₂, 200–250 μ M each unlabeled deoxyribonucleoside triphosphate (dNTP), omitting those corresponding to the labeled nucleotide(s), 50 mM Tris-

¹ P. W. J. Rigby, M. Dieckmann, C. Rhodes, and P. Berg, *J. Mol. Biol.* 113, 237 (1977).

² J. Meinkoth and G. Wahl, *Anal. Biochem.* 138, 267 (1984).

³ G. M. Wahl, M. Stern, and G. R. Stark, *Proc. Natl. Acad. Sci. U.S.A.* 76, 3683 (1979).

⁴ R. H. Singer, J. B. Lawrence, and C. Villave, *BioTechniques* 4, 230 (1986).

Nucleic acid hybridisation

a practical approach

Edited by

B D Hames

Department of Biochemistry,
University of Leeds, Leeds, England

S J Higgins

Department of Biochemistry,
University of Leeds, Leeds, England

 **IRL PRESS**

Oxford · Washington DC

J.E. Arrand





Figure 2. The preparation of probes by nick translation. \uparrow , original nick position; ∇ , final nick position; \longrightarrow , labelled strand; Pol I, *E. coli* DNA polymerase I; dNTPs, deoxyribonucleoside triphosphates; ds and ss DNA, double- and single-stranded DNA, respectively.

Nick translation can utilise any deoxyribonucleotide labelled with ^{32}P in the α position. $[^{125}\text{I}]$ -, $[^3\text{H}]$ - and non-radioactive biotinylated nucleotides can also be incorporated. With α - ^{32}P -labelled nucleotides, final specific activities of 5×10^8 d.p.m./ μg DNA can be achieved. The detailed protocol is as follows.



molecules. Later one of the enzymes of DNA replication must remove the RNA primers.

After DNA polymerase III has elongated the lagging strand up to the next RNA primer, the enzyme releases from the DNA. Then DNA polymerase I binds and begins excising the ribonucleotides and at the same time adding deoxynucleotides at the 3' end (Fig. 3.5). As it moves, it translates the nick in the 5'-to-3' direction. The ability to substitute radioactive deoxyribonucleotides *in vitro* for nonradioactive ones by nick translation is a convenient method for making DNA radioactive for genetic engineering experiments. When the RNA primer has been completely removed and DNA pol I has released from the DNA, DNA ligase can bind and seal the




Figure 3.5 Conversion of elongating Okazaki fragments to a completed chain by the actions of DNA pol I and DNA ligase.

Genetics and Molecular Biology

ROBERT SCHLEIF
Brandeis University



ADDISON-WESLEY PUBLISHING COMPANY

Reading, Massachusetts • Menlo Park, California • Don Mills, Ontario • Wokingham, England
Amsterdam • Sydney • Singapore • Tokyo • Mexico City • Bogotá • Santiago • San Juan